

(d) *Initial proof.* Except when the proof of spirits is used in making the gauge prescribed in 27 CFR 19.383 or in making a gauge for determination of tax, the initial determination of proof made on the bonded premises of a distilled spirits plant for such spirits may be used whenever a subsequent gauge is required to be made at that same plant provided that no material has been added to change the proof of the spirits.

(Sec. 201, Pub. L. 85-859, 72 Stat. 1358, as amended, 1362, as amended (26 U.S.C. 5204, 5211))

[T.D. ATF-198, 50 FR 8535, Mar. 1, 1985]

§ 30.32 Determination of proof obscuration.

(a) *General.* Proof obscuration of spirits containing more than 400 but not more than 600 milligrams of solids per 100 milliliters shall be determined by one of the following methods. The evaporation method may be used only for spirits in the range of 80-100 degrees at gauge proof.

(b) *Evaporation method.* Evaporate the water and alcohol from a carefully measured 25 milliliter sample of spirits, dry the residue at 100 degrees centigrade for 30 minutes and then weigh the residue precisely. Multiply the weight of the residue by 4 to determine the weight of solids in 100 milliliters. The resulting weight per 100 milliliters multiplied by 4 will give the obscuration. Experience has shown that 0.1 gram (100 milligrams) of solids per 100 milliliters of spirits in the range of 80-100 degrees proof will obscure the true proof by 0.4 of one degree of proof. For example, if the weight of solids remaining after evaporation of 25 milliliters 0.125 gram, the amount of solids present in 100 milliliters of the spirits is 0.50 gram (4 times 0.125). The obscuration is 4 times 0.50, which is two degrees of proof. This value added to the temperature corrected hydrometer reading will give the true proof.

(c) *Distillation method.* Determine the apparent proof and temperature of the sample of spirits and then distill a carefully measured sample in a small laboratory still, and collect a quantity of the distillate, 1 or 2 milliliters less than the original sample. The distillate is adjusted to the original temperature

and restored to the original volume by addition of distilled water. The proof of the restored distillate is then determined by use of a precision hydrometer and thermometer in accordance with the provisions of § 13.23 to the nearest 0.1 degree of proof. The difference between the proof so determined and the apparent proof of the undistilled sample is the obscuration; or

(d) *Pycnometer method.* Determine the specific gravity of the undistilled sample, distill and restore the samples as provided in paragraph (c) of this section and determine the specific gravity of the restored distillate by means of a pycnometer. The specific gravities so obtained will be converted to degrees of proof by interpolation of Table 6 to the nearest 0.1 degree of proof. The difference in proof so obtained is the obscuration.

(Sec. 201, Pub. L. 85-859, 72 Stat. 1358, as amended (26 U.S.C. 5204))

[T.D. ATF-198, 50 FR 8535, Mar. 1, 1985, as amended by T.D. ATF-381, 61 FR 37004, July 16, 1996]

DETERMINATION OF QUANTITY

§ 30.36 General requirements.

The quantity determination of distilled spirits that are withdrawn from bond in bulk upon tax determination or payment shall be by weight. The quantity of other distilled spirits or denatured spirits may be determined by weight or by volume. When the quantity of distilled spirits or denatured distilled spirits is determined by volume, such determination may be by meter as provided in 27 CFR Part 19, or when approved by the appropriate TTB officer, another method or device.

(Sec. 201, Pub. L. 85-859, 72 Stat. 1358, as amended (26 U.S.C. 5204))

[T.D. ATF-198, 50 FR 8535, Mar. 1, 1985]

DETERMINATION OF QUANTITY BY WEIGHT

§ 30.41 Bulk spirits.

When spirits (including denatured spirits) are to be gauged by weight in bulk quantities, the weight shall be determined by means of weighing tanks, mounted on accurate scales. Before each use, the scales shall be balanced